

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

49

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07F 7/18, C08G 73/02, C03B 19/14	A1	(11) International Publication Number: WO 00/21967 (43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/US99/23794 (22) International Filing Date: 13 October 1999 (13.10.99) (30) Priority Data: 09/172,190 13 October 1998 (13.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/172,190 (CIP) Filed on 13 October 1999 (13.10.99) (71) Applicant (for all designated States except US): AFFYMETRIX, INC. [US/US]; 3380 Central Expressway, Santa Clara, CA 95051 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MCGALL, Glenn [US/US]; 1121 Sladky Avenue, Mountain View, CA 94043 (US). FORMAN, Jonathan, Eric [US/US]; 524 Shadow Court, San Jose, CA 95129 (US). (74) Agents: JOHNSTON, Madeline, I. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: FUNCTIONALIZED SILICON COMPOUNDS, THEIR SYNTHESIS AND USE (57) Abstract <p>Provided are functionalized silicon compounds and methods for their synthesis and use. The functionalized silicon compounds include at least one activated silicon group and at least one derivatizable functional group. Exemplary derivatizable functional groups include hydroxyl, amino, carboxyl and thiol, as well as modified forms thereof, such as activated or protected forms. The functionalized silicon compounds may be covalently attached to surfaces to form functionalized surfaces which may be used in a wide range of different applications. In one embodiment, the silicon compounds are attached to the surface of a substrate comprising silica, such as a glass substrate, to provide a functionalized surface on the substrate, to which molecules, including polypeptides and nucleic acids, may be attached. In one embodiment, after covalent attachment of a functionalized silicon compound to the surface of a solid silica substrate to form a functionalized coating on the substrate, an array of nucleic acids may be covalently attached to the substrate. Thus, the method permits the formation of high density arrays of nucleic acids immobilized on a substrate, which may be used, for example, in conducting high volume nucleic acid hybridization assays.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

FUNCTIONALIZED SILICON COMPOUNDS, THEIR SYNTHESIS AND USE

5

TECHNICAL FIELD

This application relates to silicon compounds, methods of making silicon compounds, and methods for use of silicon compounds as silylating agents in the treatment
10 of surfaces, such as glass.

BACKGROUND ART

Silylating agents have been developed in the art which react with and coat surfaces,
15 such as silica surfaces. For example, silylating agents for use in modifying silica used in high performance chromatography packings have been developed. Monofunctional silylating agents have been used to form monolayer surface coatings, while di- and tri-functional silylating agents have been used to form polymerized coatings on silica surfaces. Many silylating agents, however, produce coatings with undesirable properties including
20 instability to hydrolysis and the inadequate ability to mask the silica surface which may contain residual acidic silanols.

Silylating agents have been developed for the silylation of solid substrates, such as glass substrates, that include functional groups that may be derivatized by further covalent reaction. The silylating agents have been immobilized on the surface of substrates, such as
25 glass, and used to prepare high density immobilized oligonucleotide probe arrays. For example, N-(3-(triethoxysilyl)-propyl)-4-hydroxybutyramide (PCR Inc., Gainesville, FL and Gelest, Tullytown, PA) has been used to silylate a glass substrate prior to photochemical synthesis of arrays of oligonucleotides on the substrate, as described in McGall *et al.*, *J. Am. Chem. Soc.*, 119:5081-5090 (1997), the disclosure of which is
30 incorporated herein by reference.

Hydroxyalkylsilyl compounds that have been used to prepare hydroxyalkylated substances, such as glass substrates. N,N-bis(hydroxyethyl) aminopropyl-triethoxysilane has been used to treat glass substrates to permit the synthesis of high-density

oligonucleotide arrays. McGall *et al.*, *Proc. Natl. Acad. Sci.*, 93:13555-13560 (1996); and Pease *et al.*, *Proc. Natl. Acad. Sci.*, 91:5022-5026 (1994), the disclosures of which are incorporated herein. Acetoxypropyl-triethoxysilane has been used to treat glass substrates to prepare them for oligonucleotide array synthesis, as described in PCT WO 97/39151, the disclosure of which is incorporated herein. 3-Glycidoxy propyltrimethoxysilane has been used to treat a glass support to provide a linker for the synthesis of oligonucleotides. EP Patent Application No. 89 120696.3.

Methods have been developed in the art for stabilizing surface bonded silicon compounds. The use of sterically hindered silylating agents is described in Kirkland *et al.*, *Anal. Chem.* 61: 2-11 (1989); and Schneider *et al.*, *Synthesis*, 1027-1031 (1990). However, the use of these surface bonded silylating agents is disadvantageous, because they typically require very forcing conditions to achieve bonding to the glass, since their hindered nature makes them less reactive with the substrate.

It is an object of the invention to provide functionalized silicon compounds that are provided with derivatizable functional groups, that can be used to form functionalized coatings on materials, such as glass. It is a further object of the invention to provide functionalized silicon compounds that can be used to form coatings on materials that are stable under the conditions of use.

DISCLOSURE OF THE INVENTION

Provided are functionalized silicon compounds and methods for their use. The functionalized silicon compounds include an activated silicon group and a derivatizable functional group. Exemplary derivatizable functional groups include hydroxyl, amino, carboxyl and thiol, as well as modified forms thereof, such as activated or protected forms. The functionalized silicon compounds may be covalently attached to surfaces to form functionalized surfaces which may be used in a wide range of different applications. In one embodiment, the silicon compounds are attached to the surface of a substrate comprising silica, such as a glass substrate, to provide a functionalized surface on the silica containing substrate, to which molecules, including polypeptides and nucleic acids, may be attached. In one preferred embodiment, after covalent attachment of a functionalized silicon

compound to the surface of a solid silica substrate to form a functionalized coating on the substrate, an array of nucleic acids may be covalently attached to the substrate. Thus, the method permits the formation of high density arrays of nucleic acids immobilized on a substrate, which may be used in conducting high volume nucleic acid hybridization assays.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the structure of the functionalized silicon compounds VI and VII and compounds of Formula 6a.

10

Figure 2 shows the structure of the functionalized silicon compound VIII.

Figure 3 shows schemes for the synthesis of compounds IX and X.

Figure 4 is a scheme showing the synthesis of compounds of Formula 5 or 6.

Figure 5 show schemes for the synthesis of compounds XII, XIII and compounds of Formula 8.

15

Figure 6 shows schemes for the synthesis of compounds XV, VI and compounds of Formula 9.

Figure 7 shows schemes showing the synthesis of compounds of Formula 10 or 11.

Figure 8 shows schemes showing the synthesis of compounds of Formula 12 or 13.

Figure 9 is a scheme of the synthesis of compounds of Formula 16b.

20

Figure 10 illustrates the structure of compounds of Formula 14 and 15.

Figure 11 is a graph of stability of silicon compound bonded phases vs. time.

Figure 12 is a graph of hybridization fluorescence intensity vs. silane.

Figure 13 is a scheme showing the synthesis of silicon compounds XVIa-e.

Figure 14 is a scheme showing the synthesis of silicon compounds XVIIa-f.

25

Figure 15 shows the structure of compounds of the general Formulas 17-21.

Figure 16 shows the structure of some exemplary silicon compounds.

Figure 17 shows another embodiment of exemplary silicon compounds XXI and XXII.

XXII.

Figure 18 shows the structure of exemplary silicon compounds XXIII, XXV and XXVI.

30

XXVI.

Figure 19 shows the structure of exemplary silicon compounds XXIX and XXX.

Figure 20 shows the structure of exemplary silicon compounds XXIa-b, XXIIa-b and XXIIIa-b.

Figure 21 is a scheme showing the synthesis of silicon compounds XIX and XX.

Figure 22 is a scheme showing the synthesis of silicon compounds XXI and XXII.

5 Figure 23 is a scheme showing the synthesis of silicon compound XXIII.

Figure 24 is a scheme showing the synthesis of silicon compound XXIV.

Figure 25 is a scheme showing the synthesis of silicon compound XXVIII.

Figure 26 is a scheme showing the synthesis of silicon compounds XXVI and XXV.

Figure 27 is a scheme showing the synthesis of silicon compounds XXIX and XXX.

10 Figure 28 is a scheme showing the synthesis of silicon compounds XXXIa-b and XXXIIa-b.

Figure 29 is a scheme showing the synthesis of silicon compounds XXXIIa-b.

Figure 30 is a graph of normalized intensity vs. silane for silicon compounds bound to a solid substrate.

15 Figure 31 is a graph of normalized hybridization fluorescence intensity vs. silane.

MODES FOR CARRYING OUT THE INVENTION

20 Functionalized silicon compounds are provided, as well as methods for their synthesis and use. The functionalized silicon compounds may be used to form functionalized coatings on a variety of surfaces such as the surfaces of glass substrates.

Functionalized Silicon Compounds

A variety of functionalized silicon compounds, which are available commercially, or which may be synthesized as disclosed herein, may be used in the methods disclosed herein to react with surfaces to form functionalized surfaces which may be used in a wide
5 range of different applications. In one embodiment, the functionalized silicon compounds are covalently attached to surfaces to produce functionalized surfaces on substrates. For example, the silicon compounds may be attached to the surfaces of glass substrates, to provide a functionalized surface to which molecules, including polypeptides and nucleic acids, may be attached.

10 As used herein, the term "silicon compound" refers to a compound comprising a silicon atom. In a preferred embodiment, the silicon compound is a silylating agent comprising an activated silicon group, wherein the activated silicon group comprises a silicon atom covalently linked to at least one reactive group, such as an alkoxy or halide, such that the silicon group is capable of reacting with a functional group, for example on a
15 surface of a substrate, to form a covalent bond with the surface. For example, the activated silicon group on the silicon compound can react with the surface of a silica substrate comprising surface Si-OH groups to create siloxane bonds between the silicon compound and the silica substrate. Exemplary activated silicon groups include $-\text{Si}(\text{OMe})_3$; $-\text{SiMe}(\text{OMe})_2$; $-\text{SiMeCl}_2$; $\text{SiMe}(\text{OEt})_2$; SiCl_3 and $-\text{Si}(\text{OEt})_3$.

20 As used herein, the term "functionalized silicon compound" refers to a silicon compound comprising a silicon atom and a derivatizable functional group. In a preferred embodiment, the functionalized silicon compound is a functionalized silylating agent and includes an activated silicon group and a derivatizable functional group. As used herein, the term "derivatizable functional group" refers to a functional group that is capable of
25 reacting to permit the formation of a covalent bond between the silicon compound and another substance, such as a polymer. Exemplary derivatizable functional groups include hydroxyl, amino, carboxy, thiol, and amide, as well as modified forms thereof, such as activated or protected forms. Derivatizable functional groups also include substitutable leaving groups such as halo or sulfonate. In one preferred embodiment, the derivatizable
30 functional group is a group, such as a hydroxyl group, that is capable of reacting with activated nucleotides to permit nucleic acid synthesis. For example, the functionalized

silicon compound may be covalently attached to the surface of a substrate, such as glass, and then derivatizable hydroxyl groups on the silicon compound may be reacted with an activated phosphate group on a protected nucleotide phosphoramidite or H-phosphonate, and then stepwise addition of further protected nucleotide phosphoramidites or H-phosphonates can result in the formation of a nucleic acid covalently attached to the support. The nucleic acids also may be attached to the derivatizable group via a linker. In a further embodiment, arrays of nucleic acids may be formed covalently attached to the substrate which are useful in conducting nucleic acid hybridization assays.

The term "polynucleotide" or "nucleic acid" as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides, that comprise purine and pyrimidine bases, or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. The backbone of the polynucleotide can comprise sugars and phosphate groups, as may typically be found in RNA or DNA, or modified or substituted sugar or phosphate groups. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-nucleotide components.

The functionalized silicon compounds used to form coatings on a surface may be selected, and obtained commercially, or made synthetically, depending on their properties under the conditions of intended use. For example, functionalized silicon compounds may be selected for silanization of a substrate that are stable after the silylation reaction to hydrolysis.


For example, in one embodiment, the functionalized silicon compounds are used to form a coating on a solid substrate, and include functional groups that permit the covalent attachment or synthesis of nucleic acid arrays to the solid substrate, such as glass. The resulting substrates are useful in nucleic acid hybridization assays, which are conducted, for example in aqueous buffers. In one embodiment, preferred are silicon compounds that produce coatings that are substantially stable to hybridization assay conditions, such as phosphate or TRIS buffer at about pH 6-9, and at elevated temperatures, for example, about 25 - 65°C, for about 1 to 72 hours, such that hydrolysis is less than about 90%, *e.g.*, less than about 50%, or *e.g.*, less than about 20%, or about 10%. The functionalized surfaces on the substrate, formed by covalent attachment of functionalized silicon compounds,

advantageously are substantially stable to provide a support for biomolecule array synthesis and to be used under rigorous assay conditions, such as nucleic acid hybridization assay conditions.

5 The functionalized silicon compound in one embodiment includes at least one activated silicon group and at least one derivatizable functional group. In one embodiment, the functionalized silicon compound includes at least one activated silicon group and a plurality of derivatizable functional groups, for example, 2, 3, 4 or more derivatizable functional groups. In another embodiment, the functionalized silicon compound includes at least one derivatizable functional group and a plurality of activated silicon groups, for
10 example, 2, 3, 4 or more activated silicon groups. Methods of making the functionalized silicon compounds are provided as disclosed herein, as well as methods of use of the functionalized silicon compounds, including covalent attachment of the silicon compounds to surfaces of substrates to form functionalized surfaces, and further derivation of the surfaces to provide arrays of nucleic acids for use in assays on the surfaces.

15 In one embodiment, there is provided a method of functionalizing a surface, the method comprising covalently attaching to the surface a functionalized silicon compound, wherein the functionalized silicon compound comprises at least one derivatizable functional group and a plurality of activated silicon groups, for example, 2, 3, 4 or more activated silicon groups. The method may further comprise covalently attaching a plurality
20 of functionalized silicon compounds to the surface, and forming an array of nucleic acids covalently attached to the functionalized silicon compounds on the surface.

Exemplary functionalized silicon compounds include compounds of Formula 1 shown below:




25

Formula 1


wherein R_1 and R_2 are independently a reactive group, such as halide or alkoxy, for example $-OCH_3$ or $-OCH_2CH_3$, and R_3 is alkoxy, halide or alkyl; and wherein R_4 is a hydrophobic and/or sterically hindered group. In the functionalized silylating agents of Formula 1, R_4 may be alkyl or haloalkyl, for example, $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, $-C(CH_3)_3$, or $-CH(CF_3)_2$. A hydrophobic and/or sterically hindered R_4 group, such as isopropyl or isobutyl, may be used to increase the hydrolytic stability of the resulting surface layer. Further hydrophobicity may be imparted by the use of a fluorocarbon R_4 group, such as hexafluoroisopropyl $((CF_3)_2CH-)$.

An exemplary compound of Formula 1 is silicon compound I below:



In general, silicon compounds provide uniform and reproducible coatings. Silicon compounds with one derivatizable functional group can provide a lower concentration of surface derivatizable functional groups at maximum coverage of the substrate than the silicon compounds including multiple derivatizable functional groups. Silicon compounds with one derivatizable functional group, such as silicon compounds of Formula 1, however, which include a hydrophobic and/or sterically hindered R group, such as isopropyl or isobutyl, are advantageous since the hydrophobic or sterically hindered R group increases the hydrolytic stability of the resulting surface layer.

In one embodiment the functionalized silicon compounds of Formula 2 are provided:



Formula 2

5


In Formula 2, in one embodiment, R_1 , R_2 and R_3 are independently a reactive group, such as alkoxy or halide, for example, $-OCH_3$, or $-OCH_2CH_3$, and wherein, in one embodiment, R_1 , R_2 and R_3 are each $-OCH_3$. In one embodiment R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example $-OCH_3$ or $-OCH_2CH_3$, and R_3 is an alkoxy or halide group or an alkyl group, such as $-CH_3$, or substituted alkyl group.

10

In Formula 2, in one embodiment, L_1 and L_2 are independently alkyl, preferably $-(CH_2)_n-$, wherein $n = 2$ to 10 , *e.g.*, 3 to 4, or *e.g.*, 2-3.

15

In Formula 2, in one embodiment, A_1 is H or a moiety comprising one or more derivatizable functional groups. In one embodiment, A_1 is a moiety comprising an amino group or a hydroxyl group, such as $-CH_2CH_2OH$. In another embodiment, A_1 is, for example, a branched hydrocarbon including a plurality of derivatizable functional groups, such as hydroxyl groups. In one embodiment in Formula 2, A_1 is:




20

wherein p is 1-10, q is 0 or 1, and r is 2-5.


In one embodiment of Formula 2, R_1 and R_2 are independently alkoxy or halide; R_3 is alkoxy, halide or alkyl; L_1 and L_2 are both $-(CH_2)_n-$, wherein $n = 2$ to 10 , *e.g.*, 2 to 3 ; and A_1 is H or a moiety comprising one or more derivatizable functional groups.

Exemplary compounds of Formula 2 include compounds II, III and IV below.


5 Other silicon compounds of Formula 2 that may be used to form functional surface coatings with enhanced hydrolytic stability include silicon compounds IX and X, shown in Figure 3. In compound VIII, the triethoxysilyl group is shown by way of example, however alternatively, the activated silicon group may be other activated silicon groups or mixtures thereof, such as trimethoxysilyl. In another embodiment, there is provided a
 10 compound of Formula 11, wherein n is, for example, 1 to 10 , *e.g.*, $1-3$, and G is a derivatizable functional group, such as hydroxyl, protected hydroxyl or halide such as Cl or Br, as shown in Figure 7.



II



III



IV

20 In another embodiment, silicon compounds of Formula 14 in Figure 10 are provided wherein, R_1 , R_2 are independently a reactive group such as alkoxy, for example $-OCH_3$ or $-OCH_2CH_3$, or halide; and R_3 is a reactive group such as alkoxy or halide, or optionally alkyl.

Another embodiment of Formula 2 is as follows:

In one embodiment of Formula 2, R_1 , R_2 , R_3 are independently a reactive group, such as alkoxy or halide, for example, $-\text{OCH}_3$, or $-\text{OCH}_2\text{CH}_3$, for example, in one embodiment, R_1 , R_2 and R_3 are each $-\text{OCH}_3$; or in another embodiment, R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example $-\text{OCH}_3$ or $-\text{OCH}_2\text{CH}_3$, and R_3 is an alkoxy or halide group or an alkyl group, such as $-\text{CH}_3$, or substituted alkyl group.

In one embodiment of Formula 2, L_1 and L_2 are independently alkyl, for example, linear or branched alkyl or heteroalkyl, *e.g.*, C1-C25 alkyl, for example, $-(\text{CH}_2)_n-$, wherein $n = 2$ to 10 , *e.g.*, 3 to 4 , or *e.g.*, $2-3$. For example, L_1 and L_2 may optionally comprise a heteroalkyl comprising a heteroatom such as O, S, or N. Each L_1 and L_2 independently comprise one or more derivatizable groups, *e.g.*, 1-4 derivatizable groups, such as hydroxyl, amino or amido.

In Formula 2, in one embodiment, A_1 is H or a moiety comprising one or more derivatizable functional groups. In one embodiment, A_1 is a moiety comprising an amino group or a hydroxyl group, such as $-\text{CH}_2\text{CH}_2\text{OH}$. In another embodiment, A_1 is, for example, a linear or branched alkyl or heteroalkyl group including a plurality of derivatizable functional groups, for example, 1, 2, or 3 derivatizable groups. In one embodiment, A_1 may comprise a linear or branched alkyl or heteroalkyl, wherein one or more carbon atoms of the alkyl group is functionalized, for example, to comprise an amide.

Examples of compounds include compounds XVIa-e shown in Figure 13, and compounds XVIIIa-f shown in Figure 14. Other examples include compound XII in Figure 5 and compound XV in Figure 6, as well as compounds XXIX and XXX in Figure 19.

In a further embodiment, compounds of Formulas 17, 18, 19, and 20 shown in Figure 15 are provided.

In one embodiment of the compounds of Formulas 17- 20, R_1 , R_2 , R_3 are independently a reactive group, such as alkoxy or halide, for example, $-\text{OCH}_3$, or $-\text{OCH}_2\text{CH}_3$, for example, in one embodiment, R_1 , R_2 and R_3 are each $-\text{OCH}_3$; or in

another embodiment, R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example $-OCH_3$ or $-OCH_2CH_3$, and R_3 is an alkoxy or halide group or an alkyl group, such as $-CH_3$, or substituted alkyl group.

5 In one embodiment of the compounds of Formulas 17-20, L_1 , L_2 , and L_3 are independently linear or branched alkyl or heteroalkyl, *e.g.*, C1-25 alkyl, for example, $-(CH_2)_n-$, wherein $n = 2$ to 10 , *e.g.*, 3 to 4 , or *e.g.*, $2-3$. For example, L_1 , L_2 , and L_3 may optionally comprise a heteroalkyl comprising a heteroatom such as O, S, or N. Each L_1 , L_2 , and L_3 independently optionally comprise one or more
10 derivativizable groups, *e.g.*, 1-4 derivativizable groups, such as hydroxyl or an amino group.

In one embodiment of Formulas 17-20, A_1 and A_2 may independently comprise H or a moiety comprising one or more derivativizable functional groups. In one embodiment, A_1 and A_2 are independently moieties comprising an amino group or a hydroxyl group, such as $-CH_2CH_2OH$. In another embodiment, A_1 and A_2 may
15 independently comprise, for example, a linear or branched alkyl or heteroalkyl including a plurality of derivativizable functional groups, for example, 1, 2, or 3 derivativizable groups. In one embodiment, A_1 and A_2 may independently comprise a linear or branched alkyl or heteroalkyl, wherein one or more carbon atoms of the alkyl group is functionalized, for example, to an amide.

20 In one embodiment of the compounds of Formulas 17-20, B_1 and B_2 are independently a branching group, for example alkyl, a heteroatom, or heteroalkyl, for example a C1-12 alkyl.

In one embodiment of the compounds of Formulas 17-20, L_4 is a direct bond or a linker, for example, C1-12 alkyl or heteroalkyl optionally comprising one more
25 derivativizable groups.

R in one embodiment is H or alkyl or heteroalkyl, for example C1-12 alkyl, or in another embodiment is acyl, for example $HC(=CH_2)CO-$ or $MeC(=CH_2)CO-$.

30 Examples of compounds of Formula 17 include compounds XIX, XX and XXIV in Figure 16. Examples of compounds of Formula 19 include compounds XXI and XXII in

Figure 17. Other examples of compounds include compounds XXVII and XXVIII in Figure 16.

In one embodiment, compounds of Formula 18 include compounds XXXIa and XXXIb shown in Figure 20. Compounds of Formula 20 include compounds XXXIIa and XXXIIb shown in Figure 20.

In another embodiment, the compounds may include a single silicon group, as for example compounds XXXIIIa and XXXIIIb shown in Figure 20.

In another embodiment, compounds of Formula 21 in Figure 15 are provided, wherein:

In one embodiment of Formula 21, R_1 , R_2 , R_3 are independently a reactive group, such as alkoxy or halide, for example, $-OCH_3$, or $-OCH_2CH_3$, for example, in one embodiment, R_1 , R_2 and R_3 are each $-OCH_3$; or in another embodiment, R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example $-OCH_3$ or $-OCH_2CH_3$, and R_3 is an alkoxy or halide group or an alkyl group, such as $-CH_3$, or substituted alkyl group.

In one embodiment of Formula 21, L_1 , L_2 , L_3 , and L_4 are independently a direct bond, or linear or branched alkyl or heteroalkyl, *e.g.*, C1-25 alkyl, for example, $-(CH_2)_n$, wherein $n = 2$ to 10 , *e.g.*, 3 to 4 , or *e.g.*, $2-3$. For example, L_1 , L_2 , L_3 and L_4 may optionally comprise a heteroalkyl comprising a heteroatom such as O, S, or N. Each L_1 , L_2 , L_3 and L_4 independently optionally comprise one or more derivatizable groups, *e.g.*, 1-4 derivatizable groups, such as hydroxyl or an amino group.

In one embodiment, B_1 is a branching group, for example alkyl, a heteroatom, or heteroalkyl, for example a C1-12 alkyl.

In one embodiment, A_1 , A_2 , A_3 , and A_4 may independently comprise H or a moiety comprising one or more derivatizable functional groups. In one embodiment, A_1 , A_2 , A_3 , and A_4 are independently a moiety comprising an amino group or a hydroxyl group, such as $-CH_2CH_2OH$. In another embodiment, A_1 , A_2 , A_3 , and A_4 may independently comprise, for example, an alkyl, such as a linear or branched alkyl, including a plurality of derivatizable functional groups, for


example, 1, 2 or 3 derivatizable groups. In one embodiment, A_1 , A_2 , A_3 , and A_4 may independently comprise a linear or branched alkyl or heteroalkyl, wherein one or more carbon atoms of the alkyl group is functionalized, for example, to comprise an amide.

5

Embodiments of compounds of Formula 21 include compounds XXIII, XXV and XXVI shown in Figure 18.

In a further embodiment, compounds of Formula 3 are provided:

10



Formula 3

15

In Formula 3, in one embodiment, R_1 , R_2 and R_3 are independently reactive groups, such as alkoxy or halide, for example, $-OCH_3$, or $-OCH_2CH_3$, and wherein, in one embodiment, R_1 , R_2 and R_3 are each $-OCH_3$. In one embodiment R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example $-OCH_3$ or $-OCH_2CH_3$, and R_3 is an alkoxy or halide group or an alkyl group, such as $-CH_3$, or substituted alkyl group.

20

In Formula 3, in one embodiment, L_1 , L_2 , and L_3 are independently a linker, for example, a straight chain saturated hydrocarbon, such as $-(CH_2)_n-$, wherein $n = 1$ to 10, or 1 to 5, or, *e.g.*, 2 to 3.

25

In Formula 3, in one embodiment, A_1 and A_2 are independently H or moieties comprising one or more derivatizable functional groups,


such as hydroxyl or amino groups, or modified forms thereof, such as protected forms. In another embodiment, A₁ and A₂ each comprise a plurality of derivativizable functional groups. For example, A₁ and A₂ may each comprise a branched moiety including a plurality of derivativizable functional groups, such as hydroxyl groups.

In one embodiment of Formula 3, R₁ and R₂ are independently alkoxy or halide; R₃ is alkoxy, halide or alkyl; L₁, L₂, and L₃ are independently -(CH₂)_n-, wherein n is 2-10; and A₁ and A₂ are independently a moiety comprising one or more derivativizable functional groups.

In another embodiment of Formula 3, A₁ is -L₄-G₁ and A₂ is -L₅-G₂; R₁ and R₂ are independently alkoxy or halide; R₃ is alkoxy, halide or alkyl; L₁, L₂, L₃, L₄ and L₅ are -(CH₂)_n-, wherein n is 1 to 10, for example 2 to 3; and G₁ and G₂ are independently a moiety comprising one or more derivativizable functional groups. In another embodiment, L₁, L₄, and L₅ are -(CH₂)₂-, L₂ and L₃ are -(CH₂)₃-, and G₁ and G₂ are -OH.

In another embodiment, silicon compounds of Formula 15 in Figure 10 are provided, wherein R₁, R₂ are independently alkoxy, for example -OCH₃ or -OCH₂CH₃, or halide; and R₃ is alkoxy, alkyl, or halide.

In a further embodiment, compounds of Formula 6a in Figure 1 are provided, wherein n is 1-3, for example 2 or 3. Exemplary functionalized silicon compounds include compound V below, and compound VI shown in Figure 1.



V


Another embodiment is illustrated in Figure 7, which shows a compound of Formula 10, wherein $n = 1$ to 10, *e.g.*, 1-3, and G is a derivatizable functional group, such as hydroxyl, protected hydroxyl, or halide such as Cl or Br.

The hydrolytic stability of the silicon compound coating may be increased by
5 increasing the number of covalent bonds to the surface of the support. For example, silicon compounds II - V include two activated silicon groups for binding to a support surface, such as glass. A variety of functionalized silicon compounds including a plurality of activated silicon groups and derivatizable functional groups are useful to form functionalized coatings. A further example is compound VII shown in Figure 1. Another
10 example is silicon compound VIII shown in Figure 2, which can form up to three covalent bonds to the surface of a glass support. In compound VIII, the triethoxysilyl group is shown by way of example, however alternatively, the activated silicon group may be other activated silicon groups or mixtures thereof, such as trimethoxysilyl. Similarly, in all of the silicon compounds disclosed herein in which a representative activated silicon group, such
15 as trimethoxysilyl, is substituted on the compound, the compounds in other embodiments also may be substituted with other activated silicon groups known in the art and disclosed herein.

The silicon compounds II-VIII having multiple silicon groups enhance potentially by twice as much, or more, the hydrolytic stability in comparison to silicon compounds
20 comprising only a single silicon group, since they possess more trialkoxysilyl groups that can react, and form bonds with, a surface. The number of silicon groups in the silicon compound may be modified for different applications, to increase or decrease the number of bonds to a support such as a glass support. Silicon compounds may be used that form optimally stable surface-bonded films on glass via covalent siloxane bonds. Additionally,
25 the number of derivatizable functional groups may be increased or decreased for different applications, as illustrated by silicon compounds II-VIII. Silicon compounds may be selected for use that provide the desired optimum density of surface derivatizable groups, such as hydroxyalkyl groups, for a desired application, such as the synthesis of nucleic acid arrays, or for the optimum stability during use of the array in different applications.

30 Other embodiments of functionalized silicon compounds include compound XIII shown in Figure 5.

In another embodiment, polymeric functionalized silicon compounds of Formula 4 are provided:



Formula 4

In Formula 4, in one embodiment, x, y and z are independently 1-3 and, in one embodiment, x, y and z are each 2.

In Formula 4, in one embodiment, L₁, L₂ and L₃ are independently linkers, for example, straight chain hydrocarbons, and preferably -(CH₂)_m-, wherein m= 1-10, e.g., 2-3.

In Formula 4, in one embodiment, at least one of A, B and C is -SiR₁R₂R₃, wherein R₁ and R₂ are independently a reactive group, such as alkoxy or halide, for example, -OCH₃, or -OCH₂CH₃ and R₃ is alkoxy, halide or alkyl; and wherein the remainder of A, B and C are independently moieties comprising one or more derivatizable functional groups, such as hydroxyl groups, or amino groups, or modified forms thereof, such as protected forms, for example -OH or a branched molecule comprising one or more hydroxyl groups.

In Formula 4, in one embodiment, n is, for example, about 10 to 10,000, or, for example, about 1,000 to 10,000.

In one embodiment of Formula 4, B is -SiR₁R₂R₃, wherein R₁, R₂ and R₃ are independently alkoxy, halide or alkyl; x, y, and z are independently 2-3; L₁, L₂ and L₃ are independently -(CH₂)_m-, wherein m is 2-3; A and C are independently moieties comprising derivatizable functional groups; and n is about 10 to 10,000.

In one embodiment of Formula 4, B is $-\text{Si}(\text{OCH}_3)_3$; x, y, and z are 2; L_1 and L_3 are $-(\text{CH}_2)_2-$; L_2 is $-(\text{CH}_2)_3-$; A and C are moieties comprising derivatizable functional groups; and n is about 10 to 10,000.

Other embodiments of a polymeric functionalized silicon compound include
5 compounds of Formula 5 and 6 shown in Figure 4, wherein m is about 0 to 10, e.g., about 1 to 5, and n is about 10 to 10,000. In Formulas 5 and 6, R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example, $-\text{OCH}_3$ or $-\text{OCH}_2\text{CH}_3$, and R_3 is a reactive group, such as alkoxy or halide, or optionally alkyl, for example $-\text{CH}_3$.

Other embodiments include compounds of Formula 7 and 8, shown in Figure 5, and
10 Formula 9, shown in Figure 6, wherein n is about 10 to 10,000. In Formulas 7, 8 and 9, R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example, $-\text{OCH}_3$ or $-\text{OCH}_2\text{CH}_3$, and R_3 is a reactive group such as alkoxy or halide or optionally alkyl, for example $-\text{CH}_3$.

Further embodiments include compounds of Formula 12 and 13, shown in Figure 8,
15 wherein m is about 10 to 10,000, and n is about 1 to 10, e.g., about 5 to 10. In Formulas 12 and 13, R_1 and R_2 are independently a reactive group, such as alkoxy or halide, for example, $-\text{OCH}_3$ or $-\text{OCH}_2\text{CH}_3$, and R_3 is a reactive group, such as alkoxy or halide, or optionally alkyl, for example $-\text{CH}_3$. In Formula 12, G is a substitutable leaving group, such as hydroxy, protected hydroxy, or halo, such as $-\text{Cl}$ or $-\text{Br}$.

20 The use of a polymer permits the formation of stable films on surfaces, such as glass, due to the very large number of siloxane bonds that can be formed with the surface. The number of alkoxy silicon groups relative to the number of hydroxyalkyl groups can be selected to provide the desired density of reactive hydroxyl groups.


Synthesis of Functionalized Silicon Compounds

25 Functionalized silicon compounds for use in the methods described herein are available commercially, or may be synthesized from commercially available starting materials. Commercially available silicon compounds and a review of silicon compounds is provided in Arkles, Ed., "Silicon, Germanium, Tin and Lead Compounds, Metal Alkoxides, Diketonates and Carboxylates, A Survey of Properties and Chemistry," Gelest,
30 Inc., Tullytown, PA, 1995, the disclosure of which is incorporated herein. Functionalized silicon compounds may be synthesized using methods available in the art of organic

chemistry, for example, as described in March, *Advanced Organic Chemistry*, John Wiley & Sons, New York, 1985, and in R.C. Larock, *Comprehensive Organic Transformations*, Wiley-VCH, New York, 1989.

Methods of synthesizing compounds of Formula 1 are shown in Scheme I below.


5 Commercially available reagents which may be used in syntheses in accordance with Scheme I include 3-chloro-1-triethoxysilylpropane and ethylene oxide (Aldrich®, Milwaukee, WI).



Formula 1

Scheme I

15 A method for the conversion of bis(trimethoxysilylpropyl)amine, XIV, which is commercially available from Gelest, Inc. (Tullytown, PA) to compound II is illustrated below in Scheme II.




XIV

II


Scheme II

A method for the conversion of compound XI, bis[3-trimethoxysilyl]propyl]-ethylenediamine, which is commercially available from Gelest, Inc., Tullytown, PA, to compound V is shown below in Scheme III.



Scheme III

A method for the synthesis of compound III is shown below in Scheme IV. The reagents shown in Scheme IV are commercially available from Aldrich® (Milwaukee, WI).



Scheme IV

Reaction schemes for the synthesis of functionalized silicon compounds IX and X are provided in Figure 3. Reaction schemes for the synthesis of compounds of Formulas 5 and 6 are shown in Figure 4. Polyethyleneimine is available commercially, for example, from Aldrich®. Polyamines of Formula 5, where R_1 , R_2 and R_3 are OMe (trimethoxysilylpropyl modified (polyethyleneimine)), or R_1 is Me and R_2 and R_3 are OMe

(dimethoxymethylsilylpropyl modified (polyethylenimine)) are available from Gelest (Tullytown, PA). Figure 9 shows another embodiment of a reaction scheme using commercially available reagents, wherein the compound of Formula 16a is converted to the compound of Formula 16b.

5 Reaction schemes for the synthesis of compounds XII, XIII, and compounds of Formula 8 are shown in Figure 5. Synthesis of the reagent, N,N-bis(2-hydroxyethyl)acrylamide is described in U.S. Patent No. 3,285,886 (1966), the disclosure of which is incorporated herein.

10 Reaction schemes for the synthesis of functionalized silicon compounds XV, VI and compounds of Formula 9 are shown in Figure 6. Use of the reagent N,N-bis(2-hydroxyethyl)-2-chloro-ethylamine is described in Okubo *et al.*, Deutsches Patent 2144759 (1971), the disclosure of which is incorporated herein.

15 Figure 7 illustrates reaction schemes for the synthesis of compounds of Formulas 10 and 11. The use of the reagent, 4-chlorobutanoyl chloride, is described in Njoroge *et al.*, PCT US97/15899 (1998), the disclosure of which is incorporated herein. Other reagents include lactones, such as γ -butyrolactone, δ -valerolactone, and ϵ -caprolactone (Aldrich®). Compounds XI and XIV are commercially available from Gelest.

20 Figure 8 illustrates reaction schemes for compounds of Formulas 12 and 13. In Figure 8, G is a substitutable leaving group such as halo. Reagents in addition to those discussed above that may be used in syntheses which may be conducted as shown in Figure 8 include diethanolamine and N,N-bis(2-hydroxyethyl)glycine, which are commercially available, for example, from Aldrich®.

25 Further exemplary reaction schemes are shown in Figures 13 and 14. Figures 13 and 14 illustrate examples of methods of synthesis of compounds of Formula 2, compounds XVIa-e, and XVIIa-f. Figure 13 illustrates examples of methods of synthesis where a compound containing a primary amino function reacts with two equivalents of (3-glycidoxypentyl)trimethoxysilane (available from Gelest, Inc., Tullytown, PA) to provide tertiary amine containing compounds XVIa-XVIe. Figure 14 illustrates examples of methods of synthesis where a compound containing a primary amino function reacts with a carbamoyl chloride (produced by reaction of XIV with triphosgene or other phosgene synthon) to produce the substituted urea compounds XVIIa-XVIIe.

30

Figure 21 provides an exemplary reaction scheme for compounds of Formula 17. The exemplary compounds XIX and XX are synthesized through the common intermediate 4-amino-1,6-heptadiene. This intermediate is prepared from ethylformimidate and methylmagnesium bromide (both available from Aldrich, Milwaukee, WI); the preparation is described in Barbot, F.; *Tetrahedron Lett.* 1989, 30, 185 and Barber, H. J.; *J. Chem. Soc.* 1943, 10. Hydrosilylation of 4-amino-1,6-heptadiene followed by reaction with ethylene oxide provides XIX. Acylation of 4-amino-1,6-heptadiene with 4-chlorobutyryl chloride (Aldrich, Milwaukee, WI) followed by hydrosilylation, and subsequent reaction of the resulting disilane with diethanol amine affords XX.

Figure 22 provides an exemplary reaction scheme for compounds of Formula 19. The exemplary compounds XXI and XXII are synthesized through the common intermediate 4-allyl-4-amino-1,6-heptadiene. This intermediate can be prepared, for example, from ethylformimidate and methylmagnesium bromide (both available from Aldrich, Milwaukee, WI); the preparation is described in Barbot, F., *Tetrahedron Lett.* 1989, 30, 185; and Barber, H. J., *J. Chem. Soc.* 1943, 101. The intermediate also may be prepared from triallylborane as described in Bubnov, Y. N., *et al.*, *Russian Chem. Bull.* 1996, 45, 2598. Hydrosilylation of 4-allyl-4-amino-1,6-heptadiene followed by reaction with ethylene oxide provides XXI. Acylation of 4-allyl-4-amino-1,6-heptadiene with 4-chlorobutyryl chloride (Aldrich, Milwaukee, WI) followed by hydrosilylation, and subsequent reaction of the resulting trisilane with diethanol amine affords XXII.

Figure 23 provides an exemplary reaction scheme for compounds of Formula 21, including structure XXIII. This compound is prepared by catalytic hydrogen reduction of 4,4-dicyano-1,6-bis(triethoxysilyl)heptane followed by reaction of the resultant diamine with ethylene oxide. The dicyano intermediate is prepared, for example, by a malonitrile synthesis of 4,4-dicyano-1,6-heptadiene followed by hydrosilylation, or by alkylation of malonitrile with 2 equivalents of a 3-halo-1-(triethoxysilyl)propane compound.

Figure 24 provides an exemplary reaction scheme for a compound of Formula 17, including structure XXIV. This compound is prepared by catalytic hydrogen reduction of 4-cyano-1,6-bis(triethoxysilyl)heptane followed by reaction of the resultant amine with ethylene oxide. The cyano intermediate is prepared starting from a malonitrile synthesis of 4,4-dicyano-1,6-heptadiene followed by conversion to 4-cyano-1,6-heptadiene by reduction

with an organotin reagent as described in Curran, D. P.; *et al.*, *Synthesis*, 1991, 107.
Hydrosilylation of this diene provides 4-cyano-1,6-bis(triethoxysilyl)heptane.

Figure 25 provides an exemplary reaction scheme for compounds of Formula 19, for example, structures XXVII and XXVIII. These compounds are prepared by reaction of appropriate primary (XXVIII) or secondary (XXVII) amines with ethyl 1,7-bis(triethoxysilyl)-4-heptanoate. The ester precursor is synthesized starting from diethyl 2,2-diallylmalonate (Aldrich, Milwaukee, WI); the diester is decarboxylated following the method of Beckwith, A. C. J.; *et al.*, *J. Chem. Soc., Perkin Trans. II*, 1975, 1726, to produce ethyl 4-hepta-1,6-dieneoate; the triethoxysilyl moieties are introduced by hydrosilylation.

Figure 26 provides an exemplary reaction scheme for compounds of Formula 21, for example, structures XXV and XXVI. These compounds are prepared by reaction of appropriate primary (XXVI) or secondary (XXV) amines with diethyl 2,2-bis(3-triethoxysilylpropyl)malonate. The diester precursor can be synthesized by hydrosilylation of diethyl 2,2-diallylmalonate (Aldrich, Milwaukee, WI).

Another embodiment of a synthesis of a compound of Formula 2 is shown in Figure 27, wherein the synthesis of compounds XXIX and XXX is shown. The general method illustrated here is to N-alkylate (XXX) or N-acylate (XXIX) bis(3-trimethoxypropyl)amine (compound XI, Gelest, Inc., Tullytown, PA) with a alkyl chain containing an ester of a carboxylic acid that can be subjected to aminolysis with dihydroxyethylamine (Aldrich, Milwaukee, WI). The alkylating agent, methyl 4-chlorobutyrate, and the acylating agent, methyl 4-chloro-4-oxobutyrate, are available from Aldrich (Milwaukee, WI).

Figure 28 provides an exemplary reaction scheme for compounds of Formula 18, for example, structures XXXIa and XXXIb. Preparation begins with the formation of 1,6-heptadiene-4-ol from reaction of ethyl formate with allylmagnesium bromide (Aldrich, Milwaukee, WI). Reaction of the alcohol with zinc chloride by the method of Reeve, W., *J. Org. Chem.* 1969, 34, 1921 affords a halo diene which after hydrosilylation can be used to alkylate dihydroxyethylamine (Aldrich, Milwaukee, WI). N-acylation with acroyl chloride (XXXIa) or methacroyl chloride (XXXIb) (Aldrich, Milwaukee, WI) following the procedure of Yokota, M., *et al.*, European patent Application 97309882.5, 1998 affords the desired compounds.

Figure 29 provides an exemplary reaction scheme for compounds of Formula 20, including structures XXXIIa and XXXIIb. Preparation begins with the formation of triallylmethanol from reaction of diethyl carbonate with allylmagnesium bromide by the method of Dreyfuss, M. P., *J. Org. Chem.* 1963, 28, 3269. Reaction of the alcohol with zinc chloride by the method of Reeve, W., *J. Org. Chem.* 1969, 34, 192 affords a halo diene which after hydrosilylation can be used to alkylate dihydroxyethylamine (Aldrich, Milwaukee, WI). N-acylation with acroyl chloride (XXXIa) or methacroyl chloride (XXXIb) (Aldrich, Milwaukee, WI) following the procedure of Yokota, M.; *et al.*, European patent Application 97309882.5, 1998 affords the desired compounds.

Figure 28 provides an exemplary reaction scheme for structures XXXIIIa and XXXIIIb, compounds which contain a single silicon atom. Preparation begins with the alkylation of dihydroxyethylamine (Aldrich, Milwaukee, WI) with an appropriate 3-halo-1-trialkoxysilylpropane reagent, followed by N-acylation with acroyl chloride (XXXIIIa) or methacroyl chloride (XXXIIIb) (Aldrich, Milwaukee, WI) following the procedure of Yokota, M., *et al.*, European patent Application 97309882.5, 1998 affords the desired compounds.

Functionalized silicon compounds within the scope of the invention that may be used to form functionalized covalent coatings on surfaces that are useful in a variety of applications and assays further include amine compounds such as compound XI, as well as reaction products formed therefrom as disclosed herein.

Applications

The methods and compositions disclosed herein may be used in a variety of applications. The functionalized silicon compounds may be covalently attached to a variety of materials, to provide derivatizable functional groups on the materials.

Exemplary materials include materials that comprise a functional group that is capable of reacting with the activated silicon group of the silicon compound. For example, the material may comprise a silica material comprising surface silanols capable of reacting with the activated silicon group to form a siloxane bond between the silicon atom on the silicon compound and the silicon atom on the surface. Thus, the functionalized silicon compounds may be attached to, for example, materials comprising silica, such as glass, chromatography material, and solid supports used for solid phase synthesis, such as nucleic

acid synthesis. The functionalized silicon compounds further may be attached to materials comprising oxides such as titanium(IV) dioxide and zirconium dioxide, aluminum oxide and indium-tin oxides, as well as nitrides, such as silicon nitride.

5 Solid substrates which may be coated by the silicon compounds include any of a variety of fixed organizational support matrices. In some embodiments, the substrate is substantially planar. In some embodiments, the substrate may be physically separated into regions, for example, with trenches, grooves, wells and the like. Examples of substrates include slides, beads and solid chips. The solid substrates may be, for example, biological, nonbiological, organic, inorganic, or a combination thereof, and may be in forms including
10 particles, strands, gels, sheets, tubing, spheres, containers, capillaries, pads, slices, films, plates, and slides depending upon the intended use.

The functionalized silicon compounds used advantageously may be selected with selected properties for a particular application. Functionalized silicon compounds may be selected which can form silicon compound surface coatings that have good stability to
15 hydrolysis. Functionalized silicon compounds may be selected which have a selected reactivity with the substrate and a selected derivatizable functional group depending on the intended use.

In one embodiment, the functionalized silicon compounds may be covalently attached to the surface of a solid substrate to provide a coating comprising derivatizable
20 functional groups on the substrate, thus permitting arrays of immobilized oligomers to be covalently attached to the substrate via covalent reaction with the derivatizable functional groups. The immobilized oligomers, such as polypeptides, or nucleic acids can be used in a variety of binding assays including biological binding assays. In one embodiment, high density arrays of immobilized nucleic acid probes may be formed on the substrate, and then
25 one or more target nucleic acids comprising different target sequences may be screened for binding to the high density array of nucleic acid probes comprising a diversity of different potentially complementary probe sequences. For example, methods for light-directed synthesis of DNA arrays on glass substrates is described in McGall *et al.*, *J. Am. Chem. Soc.*, 119:5081-5090 (1997), the disclosure of which is incorporated herein.

30 Silanation of glass substrates with the silicon compounds described herein can be conducted, for example by dip-, or spin-application with a 1% - 10% solution of silicon

compound in an aqueous or organic solvent or mixture thereof, for example in 95% EtOH, followed by thermal curing. See, for example, Arkles, *Chemtech*, 7:766-778 (1997); Leyden, Ed., "Silanes Surfaces and Interfaces, Chemically Modified Surfaces," Vol. 1, Gordon & Breach Science, 1986; and Plueddemann, E.P., Ed., "Silane Coupling Reagents", Plenum Pub. Corp., 1991, the disclosures of which are incorporated herein. Methods for
5 screening target molecules for specific binding to arrays of polymers, such as nucleic acids, immobilized on a solid substrate, are disclosed, for example, in U.S. Patent No. 5,510,270, the disclosure of which is incorporated herein. The fabrication of arrays of polymers, such as nucleic acids, on a solid substrate, and methods of use of the arrays in different assays,
10 are also described in: U.S. Patent Nos. 5,677,195, 5,624,711, 5,599,695, 5,445,934, 5,451,683, 5,424,186, 5,412,087, 5,405,783, 5,384,261, 5,252,743 and 5,143,854; PCT WO 92/10092; and U.S. Application No. 08/388,321, filed February 14, 1995, the disclosures of each of which are incorporated herein. Accessing genetic information using high density DNA arrays is further described in Chee, *Science* 274:610-614 (1996), the disclosure of
15 which is incorporated herein by reference. The combination of photolithographic and fabrication techniques allows each probe sequence to occupy a very small site on the support. The site may be as small as a few microns or even a small molecule. Such probe arrays may be of the type known as Very Large Scale Immobilized Polymer Synthesis (VLSIPS®) arrays, as described in U.S. Patent No. 5,631,734, the disclosure of which is
20 incorporated herein.

In the embodiment wherein solid phase chemistry, photolabile protecting groups and photolithography are used to create light directed spatially addressable parallel chemical synthesis of a large array of polynucleotides on the substrate, as described in U.S. Patent No. 5,527,681, the disclosure of which is incorporated herein, computer tools may
25 be used for forming arrays. For example, a computer system may be used to select nucleic acid or other polymer probes on the substrate, and design the layout of the array as described in U.S. Patent No. 5,571,639, the disclosure of which is incorporated herein.

Substrates having a surface to which arrays of polynucleotides are attached are referred to herein as "biological chips". The substrate may be, for example, silicon or
30 glass, and can have the thickness of a microscope slide or glass cover slip. Substrates that are transparent to light are useful when the assay involves optical detection, as described,

e.g., in U.S. Patent No. 5,545,531, the disclosure of which is incorporated herein. Other substrates include Langmuir Blodgett film, germanium, (poly)tetrafluorethylene, polystyrene, (poly)vinylidenedifluoride, polycarbonate, gallium arsenide, gallium phosphide, silicon oxide, silicon nitride, and combinations thereof. In one embodiment, the substrate is a flat glass or single crystal silicon surface with relief features less than about 10 Angstroms.

The surfaces on the solid substrates will usually, but not always, be composed of the same material as the substrate. Thus, the surface may comprise any number of materials, including polymers, plastics, resins, polysaccharides, silica or silica based materials, carbon, metals, inorganic glasses, membranes, or any of the above-listed substrate materials. Preferably, the surface will contain reactive groups, such as carboxyl, amino, and hydroxyl. In one embodiment, the surface is optically transparent and will have surface Si-OH functionalities such as are found on silica surfaces.

In the embodiment wherein arrays of nucleic acids are immobilized on a surface, the number of nucleic acid sequences may be selected for different applications, and may be, for example, about 100 or more, or, e.g., in some embodiments, more than 10^5 or 10^8 . In one embodiment, the surface comprises at least 100 probe nucleic acids each preferably having a different sequence, each probe contained in an area of less than about 0.1 cm^2 , or, for example, between about $1 \text{ }\mu\text{m}^2$ and $10,000 \text{ }\mu\text{m}^2$, and each probe nucleic acid having a defined sequence and location on the surface. In one embodiment, at least 1,000 different nucleic acids are provided on the surface, wherein each nucleic acid is contained within an area less than about 10^{-3} cm^2 , as described, for example, in U.S. Patent No. 5,510,270, the disclosure of which is incorporated herein.

Arrays of nucleic acids for use in gene expression monitoring are described in PCT WO 97/10365, the disclosure of which is incorporated herein. In one embodiment, arrays of nucleic acid probes are immobilized on a surface, wherein the array comprises more than 100 different nucleic acids and wherein each different nucleic acid is localized in a predetermined area of the surface, and the density of the different nucleic acids is greater than about 60 different nucleic acids per 1 cm^2 .

Arrays of nucleic acids immobilized on a surface which may be used also are described in detail in U.S. Patent No. 5,744,305, the disclosure of which is incorporated

herein. As disclosed therein, on a substrate, nucleic acids with different sequences are immobilized each in a predefined area on a surface. For example, 10, 50, 60, 100, 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , or 10^8 different monomer sequences may be provided on the substrate. The nucleic acids of a particular sequence are provided within a predefined region of a substrate, having a surface area, for example, of about 1 cm^2 to 10^{-10} cm^2 . In some
5 embodiments, the regions have areas of less than about 10^{-1} , 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} , 10^{-6} , 10^{-7} , 10^{-8} , 10^{-9} , or 10^{-10} cm^2 . For example, in one embodiment, there is provided a planar, non-porous support having at least a first surface, and a plurality of different nucleic acids attached to the first surface at a density exceeding about 400 different nucleic acids/ cm^2 ,
10 wherein each of the different nucleic acids is attached to the surface of the solid support in a different predefined region, has a different determinable sequence, and is, for example, at least 4 nucleotides in length. The nucleic acids may be, for example, about 4 to 20 nucleotides in length. The number of different nucleic acids may be, for example, 1000 or more. In the embodiment where polynucleotides of a known chemical sequence are
15 synthesized at known locations on a substrate, and binding of a complementary nucleotide is detected, and wherein a fluorescent label is detected, detection may be implemented by directing light to relatively small and precisely known locations on the substrate. For example, the substrate is placed in a microscope detection apparatus for identification of locations where binding takes place. The microscope detection apparatus includes a
20 monochromatic or polychromatic light source for directing light at the substrate, means for detecting fluoresced light from the substrate, and means for determining a location of the fluoresced light. The means for detecting light fluoresced on the substrate may in some embodiments include a photon counter. The means for determining a location of the fluoresced light may include an x/y translation table for the substrate. Translation of the
25 substrate and data collection are recorded and managed by an appropriately programmed digital computer, as described in U.S. Patent No. 5,510,270, the disclosure of which is incorporated herein.

Devices for concurrently processing multiple biological chip assays may be used as described in U.S. Patent No. 5,545,531, the disclosure of which is incorporated herein.
30 Methods and systems for detecting a labeled marker on a sample on a solid support, wherein the labeled material emits radiation at a wavelength that is different from the

excitation wavelength, which radiation is collected by collection optics and imaged onto a detector which generates an image of the sample, are disclosed in U.S. Patent No. 5,578,832, the disclosure of which is incorporated herein. These methods permit a highly sensitive and resolved image to be obtained at high speed. Methods and apparatus for detection of fluorescently labeled materials are further described in U.S. Patent Nos. 5,631,734 and 5,324,633, the disclosures of which are incorporated herein.

The methods and compositions described herein may be used in a range of applications including biomedical and genetic research and clinical diagnostics. Arrays of polymers such as nucleic acids may be screened for specific binding to a target, such as a complementary nucleotide, for example, in screening studies for determination of binding affinity and in diagnostic assays. In one embodiment, sequencing of polynucleotides can be conducted, as disclosed in U.S. Patent No. 5,547,839, the disclosure of which is incorporated herein. The nucleic acid arrays may be used in many other applications including detection of genetic diseases such as cystic fibrosis, diabetes, and acquired diseases such as cancer, as disclosed in U.S. Patent Application Ser. No. 08/143,312, the disclosure of which is incorporated herein. Genetic mutations may be detected by sequencing by hybridization. In one embodiment, genetic markers may be sequenced and mapped using Type-IIs restriction endonucleases as disclosed in U.S. Patent No. 5,710,000, the disclosure of which is incorporated herein.

Other applications include chip based genotyping, species identification and phenotypic characterization, as described in U.S. Patent Application Serial No. 08/797,812, filed February 7, 1997, and U.S. Application Serial No. 08/629,031, filed April 8, 1996, the disclosures of which are incorporated herein.

Gene expression may be monitored by hybridization of large numbers of mRNAs in parallel using high density arrays of nucleic acids in cells, such as in microorganisms such as yeast, as described in Lockhart *et al.*, *Nature Biotechnology*, 14:1675-1680 (1996), the disclosure of which is incorporated herein. Bacterial transcript imaging by hybridization of total RNA to nucleic acid arrays may be conducted as described in Saizieu *et al.*, *Nature Biotechnology*, 16:45-48 (1998), the disclosure of which is incorporated herein.

All publications cited herein are incorporated herein by reference in their entirety. The invention will be further understood by the following non-limiting examples.

EXAMPLES

Example 1:

5 Silicon compounds were obtained commercially or synthesized from commercially available starting materials. Silicon compounds N,N-bis(2-hydroxyethyl)-3-aminopropyltriethoxysilane, and N-(2-hydroxyethyl)-N-methyl-3-aminopropyltriethoxysilane (compound I) were purchased from Gelest, Inc (Tullytown, PA).

10 Silicon compounds II and V were prepared as shown in Schemes II and III. Solutions of the starting materials, bis(trimethoxysilyl-propyl)amine and bis[(3-trimethoxysilyl)propyl]ethylenediamine in methanol (62 wt%, from Gelest, Inc.) were combined with 1.1-2.2 theoretical equivalents of ethylene oxide at room temperature under a dry ice-acetone condenser. The resulting solutions were analyzed by ¹H-NMR, which
15 indicated 95% conversion in some runs to the hydroxyethylated products, and these were used without further purification. In other runs, a 60-65% conversion was obtained for compound V and also was used without further purification.

¹H-NMR(CDCl₃) data are provided below:

(II): 0.55-0.70 (br m, 4H), 1.55-1.65 (br m, 4H), 2.40-2.50 (br m, 2H), 2.55-2.60 (br
20 m, 4H), 3.50 (s, MeOH), 3.55 (d, 9H), 3.75 (t, 2H);

(V): 0.55-0.70 (br m, 4H), 1.50-1.65 (br m, 4H), 2.40-2.80 (br m, 12H), 3.50 (s, MeOH), 3.40-3.60 (m-s, ~20H), 3.75 (m, ~3-4H);

Substrates were treated by a silanation procedure as follows. Glass substrates (borosilicate float glass, soda lime or fused silica, 2"x 3" x 0.027", obtained from U.S. Precision Glass (Santa Rosa, CA) were cleaned by soaking successively in Nanostrip
25 (Cyantek, Fremont, CA) for 15 minutes, 10% aqueous NaOH/70°C for 3 minutes, and then 1% aqueous HCl for 1 minute (rinsing thoroughly with deionized water after each step). Substrates were then spin-dried for 5 minutes under a stream of nitrogen at 35°C. Silanation was carried out by soaking under gentle agitation in a freshly prepared 1-2%
30 (wt/vol) solution of the silicon compound in 95:5 ethanol-water for 15 minutes. The

substrates were rinsed thoroughly with 2-propanol, then deionized water, and finally spin-dried for 5 minutes at 90°-110°C.

5 The stability of silicon compound bonded phase was evaluated. The surface hydroxyalkylsilane sites on the resulting substrates were "stained" with fluorescein in a checkerboard pattern by first coupling a MeNPOC-HEG linker phosphoramidite, image-wise photolysis of the surface, then coupling to the photo-deprotected linker sites a 1:20 mixture of fluorescein phosphoramidite and DMT-T phosphoramidite (Amersham-Pharmacia Biotech, Piscataway, NJ), and then deprotecting the surface molecules in 1:1 ethylenediamine-ethanol for 4 hr. The steps were conducted using standard protocols, as
10 described in McGall *et al.*, *J. Am. Chem. Soc.*, 119: 5081-5090 (1997), the disclosure of which is incorporated herein.

The pattern and intensity of surface fluorescence was imaged with a scanning laser confocal fluorescence microscope, which employed excitation with a 488 nm argon ion laser beam focused to a 2 micron spot size at the substrate surface. Emitted light was
15 collected through confocal optics with a 530(+15) nm bandpass filter and detected with a PMT equipped with photon counting electronics. Output intensity values (photon counts/second) are proportional to the amount of surface-bound fluorescein, so that relative yields of free hydroxyl groups within different regions of the substrate could be determined by direct comparison of the observed surface fluorescence intensities. All intensity values
20 were corrected for nonspecific background fluorescence, taken as the surface fluorescence within the non-illuminated regions of the substrate.

The relative surface reactive site density was measured. For each silicon compound tested, the number of available surface synthesis sites achieved per unit area was estimated, relative to N,N-bis(2-hydroxyethyl)-3-aminopropyl-triethoxysilane, by comparison of the
25 observed initial surface fluorescence intensities of the various substrates immediately after deprotection in ethanolic diaminoethane.

$$\text{Site Density (\% rel.)} = \frac{\text{Intensity (silicon compound "X")} \times 100}{\text{Intensity (silicon compound II)}}$$

5 To determine the relative stability of the silicon compound coatings, substrates were gently agitated on a rotary shaker at 45°C in 5xSSPE or 6xSSPE aqueous buffer (BioWhittaker Products, Walkersville, MD) at pH 7.3. Periodically, the substrates were removed from the buffer and re-scanned to measure the amount of fluorescein remaining bound to the surface.

10 The results are shown in Figure 11. As shown in Figure 11, more of the fluorescein tag remained bound to the substrate after prolonged exposure to the aqueous buffer in the case of substrates silanated with II or V, than remained bound to the substrates silanated with N,N-bis(2-hydroxyethyl)-3-aminopropyltriethoxysilane or compound I. This demonstrates that the surface bonded phase obtained with silicon compounds II or V is much more stable towards hydrolysis than that obtained with N,N-bis(2-hydroxyethyl)-3-aminopropyl-triethoxysilane.

15 The hybridization performance of silanated substrates was evaluated. A comparison was made between substrates silanated with N,N-bis(2-hydroxyethyl)-3-aminopropyltriethoxysilane and compound V in terms of performance under typical hybridization assay conditions. A nucleic acid probe sequence (5'-GTC AAG ATG CTA CCG TTC AG-3') (SEQ. ID NO. 1) was synthesized photolithographically in a checkerboard array pattern (400 x 400 micron features) on the substrates that had been derivatized with either N,N-bis(2-hydroxyethyl)-3-aminopropyltriethoxysilane or silicon compound V. After deprotection in ethanolic diaminoethane, the arrays were hybridized with a fluorescein-labeled complementary "target" nucleic acid (5'-fluorescein-CTG AAC GGT AGC ATC TTG AC-3') (SEQ. ID NO. 2) at a concentration of 250 pM in 6xSSPE buffer (0.9 M NaCl, 60 mM NaH₂PO₄, 6 mM EDTA, pH 7.5) for 16 hours at 45°C. After cooling to room temperature, the target nucleic acid solution was removed, and the array was washed briefly with 6x SSPE buffer and then scanned on a confocal imaging system. The relative amount of bound target was determined from the fluorescence signal intensity.

25 30 The hybridization signal intensities obtained with the more stable substrates, derivatized with silicon compound V, were at least four times higher than those obtained with

substrates that were derivatized with N,N-bis(2-hydroxyethyl)-3-aminopropyltriethoxysilane. A graph of fluorescence signal intensity vs. silane is shown in Figure 12.

Example 2:

5 Silicon compound XVIb was prepared as shown in Figure 13. An oven dried 25 ml single-neck flask was charged with 1.07 g (1 ml, 0.0066 mol) N,N-bis(2-hydroxyethyl)-1,3-diaminopropane (TCI America, Portland, OR) and 5 ml anhydrous methanol (Aldrich Chemical; Milwaukee, WI). To this stirring solution was added 2.92 ml (3.12 g, 0.0132 mol) (3-glycidoxypentyl)trimethoxysilane (Gelest, Inc., Tullytown, PA) dropwise over a
10 30 minute period. The solution was allowed to stir under an Ar atmosphere until ^1H NMR indicated that the triplet signal for the $(\text{CH}_2)\text{CH}_2(\text{NH}_2)$ protons of the starting amine ($\delta = 2.86$ ppm) had shifted completely away (41 hours). Molecular ions corresponding to the starting materials were absent in the mass spectrum (positive ion mode) which was dominated by the $[\text{M}+\text{H}]^+$ peak of $m/z = 635.5$.

Example 3:

15 An alternate preparation of silicon compound XVIb was conducted as shown in Figure 13. An oven dried 25 ml single-neck flask was charged with 1.07 g (1 ml, 0.0066 mol) N,N-bis(2-hydroxyethyl)-1,3-diaminopropane (TCI America, Portland, OR) and 5 ml anhydrous methanol (Aldrich Chemical; Milwaukee, WI). To this stirring solution was
20 added 2.92 ml (3.12 g, 0.0132 mol) (3-glycidoxypentyl)trimethoxysilane (Gelest, Inc., Tullytown, PA) dropwise over a 30 minute period. The solution was brought to reflux and allowed to stir under an Ar atmosphere until ^1H NMR indicated that the triplet signal for the $(\text{CH}_2)\text{CH}_2(\text{NH}_2)$ protons of the starting amine ($\delta = 2.86$ ppm) had shifted completely away (17 hours). Molecular ions corresponding to the starting materials were absent in the
25 mass spectrum (positive ion mode), an $\text{M}+\text{H}^+$ peak of $m/z = 635.5$ was observed but it did not dominate the spectrum as seen in the room temperature preparation of Example 2.

Example 4:

30 Silicon compound XVIe was prepared as shown in Figure 13. An oven dried 25 ml single-neck flask was charged with 2.74 g (0.026 mol) tris(hydroxymethyl)-methylamine (Aldrich Chemical; Milwaukee, WI), 10 ml (10.70 g, 0.0452 mol) (3-glycidoxypentyl)trimethoxysilane (Gelest, Inc., Tullytown, PA), and 11ml anhydrous

methanol (Aldrich Chemical; Milwaukee, WI). The suspension was allowed to stir under an Ar atmosphere. After all the solids dissolved (6 days), ^{13}C NMR indicated that the singlet signal for the $\text{C}(\text{CH}_2\text{OH})_3$ carbon of the starting amine ($\delta = 55.38$ ppm) had shifted completely away. ^{13}C NMR (δ , ppm): 73.12 (CH-OH), 71.03 (C-N), 61.94 (CH_2OH), 50.62 (C-[OC]), 50.06 ([C]-C-O), 49.39 (CH_3OSi), 43.85 (N-C-[CH_2OH] $_3$), 22.34 (C-[CSi]), 4.81 (C-Si). Mass Spectrum: M^+ (m/z 593.4) and $[\text{M} - \text{CH}_3\text{OH}]^+$ (m/z 561.5).

Example 5:

A solution of 104 g triphosgene (0.102 mol) (Aldrich Chemical; Milwaukee, WI) and 1 liter anhydrous THF (Aldrich Chemical; Milwaukee, WI) was prepared under Ar in a 2-l oven-dried flask. To the stirring solution was slowly added 42.5 ml (30.8 g, 0.304 mol) triethylamine (Aldrich Chemical; Milwaukee, WI) and a white precipitate formed. The stirring suspension was cooled in ice and 100 ml N,N-bis(3-trimethoxysilylpropyl)amine (104 g, 0.304 mol) (compound XIV, Gelest, Inc., Tullytown, PA) was added dropwise over an hour. The reaction mixture was allowed to stir for 21.5 hours then filtered under an Ar blanket and the solids washed with anhydrous THF. The filtrate was concentrated under reduced pressure to an orange oil and dried under vacuum to give 116 g of product (95% crude yield). ^1H NMR (δ , ppm): 3.57 (s, CH_3OSi), 3.32 (dd, $\text{CH}_2\text{-NC=O}$, 4 H), 1.70 (q, [CH_2]- CH_2 -[CH_2], 4 H), 0.58 (t, CH_2Si , 4 H). ^{13}C NMR (δ , ppm): 149.14 (C=O), 53.53 (C-N-C=O), 52.25 (C-N-C=O), 50.66 (CH_3OSi), 21.83 (C-C-N), 20.82 (C-C-N), 6.50 (C-Si).

Example 6:

Silicon compound XVIIb was prepared as shown in Figure 14. An oven dried 25 ml single-neck flask was charged with 0.21 ml (0.15 g, 0.0015 mol) triethylamine (Aldrich Chemical; Milwaukee, WI), 0.61 g of a carbamoyl chloride prepared as per Example 5 (0.0015 mol), and 5 ml anhydrous THF (Aldrich Chemical; Milwaukee, WI). After addition of 0.20 g (0.0015 mol) diethanolamine (Aldrich Chemical; Milwaukee, WI) a precipitate formed. After 5 days of stirring under Ar at room temperature, the mixture was filtered and concentrated to a brown oil. After drying under vacuum, 0.62 g (87% crude yield) of material was obtained. The oil was dissolved in 0.6 ml methanol and stored. ^{13}C NMR (δ , ppm): 166.29 (C=O), 60.50 (C-OH), 57.28 (N-C-COH), 46.14 (C-C-N), 21.01 (C-[CSi]), 6.27 (C-Si). Mass Spectrum: $[\text{M}+\text{H}]^+$ (m/z 473) and $[\text{M} - \text{CH}_3\text{O}]^+$ (m/z 441).

Example 7:

Silicon compound XVIIIf was prepared as shown in Figure 14. An oven dried 25 ml single-neck flask was charged with 42.4 ml (30.8 g, 0.304 mol) triethylamine (Aldrich Chemical; Milwaukee, WI), 116 g of a carbamoyl chloride prepared as per Example 5 (0.304 mol), and 500 ml anhydrous THF (Aldrich Chemical; Milwaukee, WI). After addition of 1.07 g (1 ml, 0.0066 mol) N,N-Bis(2-hydroxyethyl)-1,3-diaminopropane (TCI America, Portland, OR), 500 ml anhydrous CH₂Cl₂ (Aldrich Chemical; Milwaukee, WI) was added. The mixture was stirred under Ar at room temperature, during which time an insoluble orange oil formed. After 18 hours, ¹H NMR indicated that the triplet signal for the (CH₂)CH₂(NH₂) protons of the starting amine (δ = 2.86 ppm) had shifted completely away. The mixture was decanted and the filtrate concentrated. The concentrate was combined with 400 ml anhydrous THF (Aldrich Chemical; Milwaukee, WI). The resultant suspension was filtered, concentrated under reduced pressure, and dried under vacuum to yield 145 g (95% crude yield) of a faint orange oil. The oil was dissolved in 150 ml methanol and stored. Mass Spectrum: [M+H]⁺ (m/z 530).

Example 8:

Coated substrates with XVib, XVIe, XVIIb, and XVIIIf were prepared by a modified version of the silanation procedure of Example 1. In this case, the silane content of the 95:5 ethanol:water bath was 1-2% by volume of the methanol/silane compound solutions described in Examples 2-4 and 6-7 and curing was done at 50°C for 2 minutes.

Example 9:

Evaluation of stability of the silicon bonded phase, relative surface reactive site density, and hybridization performance of substrates coated with XVib, XVIe, XVIIb, and XVIIIf was carried out in a single experimental protocol. First, MeNPOC-HEG linker phosphoramidite is coupled to the surface, then image-wise photolysis is used to pattern the surface into a series of stripes. Two stripes, 400 μ m \times 12800 μ m, were stained with fluorescein label to evaluate stability of the silicon bonded phase and relative surface reactive site density. A nucleic acid probe sequence was synthesized photolithographically into two additional stripes, 1600 μ m \times 12800 μ m. After patterning, the surface is deprotected in 1:1 ethylenediamine ethanol for 4 hours. The steps were conducted using standard protocols, as described in McGall *et al.*, *J. Am. Chem. Soc.*, 119:5081-5090

(1997), the disclosure of which is incorporated herein. The pattern and intensity of surface fluorescence from both fluorescein labeled patterns and adsorbed fluorescein labeled target oligonucleotide is imaged with a scanning fluorescence microscope as described in Example 1.


5 In the assay the surface was imaged after a period of 1 hour exposure to 5 nM target in 6X SSPE buffer (0.9 M NaCl, 60 mM NaH₂PO₄, 6 mM EDTA, pH 7.5) at 25°C and again after 16 hours at 45°C. Except as noted herein, the method employed for the assay is the same as described in Example 1. The fluorescence intensity was averaged and
10 evaluated at both the 1 hour and 16 hour time points. Relative surface reactive site density was determined by normalization of the signal to that obtained from compound I as described in Example 1. Figure 30 shows the normalized data at the 1 and 16 hour time points. The percentage of remaining fluorescence intensity is noted for the 16-hour time point for each silane coating.

15 Hybridization performance of substrates for matched probes is illustrated in Figure 31, where the 1 hour, 25°C intensities of the averaged background corrected fluorescence signals of the matched sequence stripes are shown. Signals are normalized to 1 = intensity of identical probes synthesized on N,N-bis(2-hydroxyethyl)-3-trimethoxypropylamine (compound I) surfaces:

CLAIMS

What is claimed is:

1. A functionalized silicon compound having a structure of Formula 2:



10 wherein R_1 and R_2 are independently selected from the group consisting of alkoxy and halide, and R_3 is selected from the group consisting of alkoxy, halide and alkyl;

wherein L_1 and L_2 are both $-(CH_2)_n-$, wherein $n = 2$ to 10 ; and


wherein A_1 is a moiety comprising one or more derivatizable functional groups.

- 15
2. The functionalized silicon compound of claim 1, wherein the derivatizable functional group is selected from the group consisting of hydroxyl, amino, amido, carboxyl, thio, halo and sulfonate.


- 20
3. The functionalized silicon compound of claim 2, wherein A_1 comprises a plurality of derivatizable functional groups.

4. The functionalized silicon compound of claim 2, wherein A_1 is a moiety comprising a hydroxyl group.

- 25
5. The functionalized silicon compound of claim 2, wherein the silicon compound is compound II:



- 5 6. A functionalized silicon compound having a structure of Formula 3:



Formula 3

10

wherein R_1 and R_2 are independently selected from the group consisting of alkoxy or halide, and R_3 is selected from the group consisting of alkoxy, halide and alkyl;

15

wherein L_1 , L_2 , and L_3 are independently $-(\text{CH}_2)_n-$, wherein n is 2-10; and wherein A_1 and A_2 are independently a moiety comprising one or more derivatizable functional groups or modified forms thereof.


20

7. The functionalized silicon compound of claim 6, wherein the one or more derivatizable functional groups are selected from the group consisting of hydroxyl, amino, carboxyl, thio, halo, amido and sulfonate.


8. The functionalized silicon compound of claim 7, wherein A_1 and A_2 each comprise a plurality of derivatizable functional groups.

9. The functionalized silicon compound of claim 7, wherein A_1 and A_2 each comprise a hydroxyl group.

10. The functionalized silicon compound of claim 7, wherein the functionalized silicon compound is compound V:



11. A functionalized silicon compound having a structure of Formula 4:



Formula 4

wherein B is $-\text{SiR}_1\text{R}_2\text{R}_3$, wherein R_1 , R_2 and R_3 are independently alkoxy, halide or alkyl;


wherein x, y, and z are independently 2 to 3;

wherein L_1 , L_2 and L_3 are independently $-(\text{CH}_2)_m-$, wherein m is 2-10;

wherein A and C are independently moieties comprising derivatizable functional groups; and

wherein n is about 10 to 10,000.

12. A functionalized silicon compound of Formula 6:




5

Formula 6

wherein R₁, R₂ and R₃ are independently alkoxy, halide or alkyl; and
 wherein *m* is about 1 to 5, and *n* is about 10 to 10,000.

10

13. A functionalized silicon compound of Formula 2








Formula 2









wherein:


- R₁ and R₂ are independently selected from the group consisting of alkoxy or halide,
 and R₃ is selected from the group consisting of alkoxy, halide and alkyl;
 L₁ and L₂ are independently alkyl or heteroalkyl optionally comprising one or more
 derivatizable groups;
 A₁ is H or a moiety comprising one or more derivatizable functional groups.

14. The silicon compound of claim 13, wherein the derivatizable functional groups are independently selected from the group consisting of hydroxyl, amino, carboxyl, thio, halo and sulfonate.





5 15. The silicon compound of claim 13 selected from the group consisting of







16. A functionalized silicon compound of Formulas 17, 18, 19, or 20:



5

wherein:

R_1 and R_2 are independently alkoxy or halide, and R_3 is alkoxy, halide or alkyl;

L_1 , L_2 and L_3 are independently alkyl or heteroalkyl, optionally comprising one or more derivatizable groups;

A₁ and A₂ independently are H or a moiety comprising one or more derivatizable functional groups;


B₁ and B₂ are independently alkyl, a heteroatom, or heteroalkyl;

L₄ is a direct bond, alkyl or heteroalkyl optionally comprising one or more
5 derivatizable groups; and


R is H, alkyl, heteroalkyl, or acyl.

17. The silicon compound of claim 16, wherein the derivatizable functional
10 groups are independently selected from the group consisting of hydroxyl, amino, carboxyl, thio, halo, amido and sulfonate.


18. The silicon compound of claim 16, wherein the compound is selected from the group consisting of




XIX



XX




XXI




XXII

15




XXXIa R' = H
XXXIb R' = CH₃

and



XXXIIa R' = H
XXXIIb R' = CH₃

19. A functionalized silicon compound of Formula 21:



Formula 21

wherein:

R₁ and R₂ are independently selected from the group consisting of alkoxy or halide, and R₃ is selected from the group consisting of alkoxy, halide and alkyl;


L₁, L₂, L₃, and L₄ are independently alkyl or heteroalkyl optionally comprising one or more derivatizable groups;

B₁ is alkyl, a heteroatom, or heteroalkyl;


A₁, A₂, A₃, and A₄ are independently H or a moiety comprising one or more derivatizable functional groups.

20. The silicon compound of claim 19, wherein the derivatizable functional groups are independently selected from the group consisting of hydroxyl, amino, carboxyl, thio, halo amido and sulfonate.

5 21. A silicon compound of claim 19, selected from the group consisting of




XXIII



XXV

and




XXVI

10 22. A method of functionalizing a surface, the method comprising covalently attaching to the surface a functionalized silicon compound, wherein the functionalized silicon compound comprises at least one derivatizable functional group and a plurality of activated silicon groups.

15 23. The method of claim 22, wherein the method comprises covalently attaching a plurality of functionalized silicon compounds to the surface; and forming an array of nucleic acids covalently attached to the functionalized silicon compounds on the surface.

24. The method of claim 22, wherein the silicon compound is a compound having a structure of Formula 2:



Formula 2

wherein R_1 and R_2 are independently selected from the group consisting of alkoxy and halide, and R_3 is selected from the group consisting of alkoxy, halide and alkyl;


wherein L_1 and L_2 are both $-(\text{CH}_2)_n-$, wherein $n = 2$ to 10 ; and

wherein A_1 is H or a moiety comprising one or more derivatizable functional groups.

25. The method of claim 24, wherein the derivatizable functional group is a hydroxyl group or modified form thereof.


26. The method of claim 24, wherein A_1 comprises a plurality of derivatizable functional groups.

27. The method of claim 24, wherein the silicon compound is compound II:



II

28. The method of claim 22, wherein the method comprising covalently attaching to the surface a functionalized silicon compound having a structure of Formula 3:



Formula 3

wherein R_1 and R_2 are independently alkoxy or halide, and R_3 is selected from the group consisting of alkoxy, halide and alkyl;


wherein L_1 , L_2 , and L_3 are $-(CH_2)_n-$, wherein n is 2 to 10; and

wherein A_1 and A_2 are independently H or a moiety comprising one or more derivatizable functional groups.

29. The method of claim 28, wherein the one or more derivatizable functional groups are hydroxyl groups or modified forms thereof.

30. The method of claim 28, wherein A_1 and A_2 each comprise a plurality of derivatizable functional groups.


31. The method of claim 28, wherein the functionalized silicon compound is compound V:



V

32. The method of claim 22, wherein the method comprises covalently attaching to the surface a functionalized silicon compound having a structure of Formula 4:

5



Formula 4

wherein B is $-\text{SiR}_1\text{R}_2\text{R}_3$, wherein R_1 , R_2 and R_3 are independently alkoxy, halide or alkyl;
 wherein x, y, and z are independently 2 to 3;
 wherein L_1 , L_2 and L_3 are independently $-(\text{CH}_2)_m-$, wherein m is 2 to 10;
 wherein A and C are independently moieties comprising derivatizable functional groups; and
 wherein n is about 10 to 10,000.

33. A method of functionalizing a surface, the method comprising covalently attaching to the surface a functionalized silicon compound according to claim 12.


34. A method of functionalizing a surface, the method comprising covalently attaching to the surface a functionalized silicon compound of claim 13.

35. The method of claim 34, wherein the method comprises covalently attaching a plurality of functionalized silicon compounds to the surface; and


forming an array of nucleic acids covalently attached to the functionalized silicon compounds on the surface.

36. The method of claim 34, wherein the derivatizable groups are hydroxyl groups or modified forms thereof.


5 37. The method of claim 34, wherein the compound is selected from the group consisting of




XVIa




XVIb



XVIc





38. A method of functionalizing a surface, the method comprising covalently attaching to the surface a functionalized silicon compound of claim 16.

5


39. The method of claim 38, wherein the method comprises covalently attaching a plurality of functionalized silicon compounds to the surface; and forming an array of nucleic acids covalently attached to the functionalized silicon compounds on the surface.

10


40. The method of claim 38, wherein the derivatizable groups are hydroxyl groups or modified forms thereof.

15


41. The method of claim 38, wherein the compound is selected from the group consisting of




XIX




XX



XXI




XXII



XXXIa R' = H
XXXIb R' = CH₃

and



XXXIIa R' = H
XXXIIb R' = CH₃


5 42. A method of functionalizing a surface, the method comprising covalently attaching to the surface a functionalized silicon compound of claim 19.

43. The method of claim 42, wherein the method comprises covalently attaching a plurality of functionalized silicon compounds to the surface; and


forming an array of nucleic acids covalently attached to the functionalized silicon compounds on the surface.

44. The method of claim 42, wherein the derivatizable groups are hydroxyl groups or modified forms thereof.

45. The method of claim 42, wherein the compound is selected from the group consisting of




XXIII



XXV

and



XXVI

46. The method of claim 22, wherein the surface is the surface of a substrate comprising silica.

47. The method of claim 24, wherein the surface is the surface of a substrate comprising silica.

48. The method of claim 28, wherein the surface is the surface of a substrate comprising silica.


49. The method of claim 32 wherein the surface is the surface of a substrate comprising silica.

50. The method of claim 34, wherein the surface is the surface of a substrate comprising silica.


51. The method of claim 38, wherein the surface is the surface of a substrate comprising silica.

52. The method of claim 42, wherein the surface is the surface of a substrate comprising silica.


53. The method of claim 22, wherein the substrate is in a form selected from the group consisting of particles, films and chips.



VI




Formula 6a



VII

FIGURE 1



VIII

FIGURE 2






FIGURE 3



Formula 5



Formula 6

FIGURE 4




FIGURE 5




FIGURE 6

7/31





FIGURE 7





FIGURE 8


9/31




Formula 16a

Formula 16b

FIGURE 9



Formula 14



Formula 15

FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13







FIGURE 14




Formula 17




Formula 18



Formula 19




Formula 20




Formula 21

FIGURE 15


16/31




XIX




XX



XXIV




XXVII




XXVIII

FIGURE 16




XXI




XXII

FIGURE 17


18/31



XXIII




XXV




XXVI

FIGURE 18




XXIX




XXX


FIGURE 19




XXXIIIa




XXXXIa




XXXIIa



XXXIIIb



XXXXIb



XXXIIb

FIGURE 20




FIGURE 21




FIGURE 22




FIGURE 23




FIGURE 24

25/31




FIGURE 25




FIGURE 26




FIGURE 27




FIGURE 28




FIGURE 29




Figure 30




Figure 31

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/23794

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07F/18 C08G73/02 C03B19/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07F C08G C03B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 393 353 A (BISHOP, C.V.) 28 February 1995 (1995-02-28) column 5, line 10-12	1,2
X	US 4 990 641 A (KABETA, K.) 5 February 1991 (1991-02-05) example 5	1,2
X	EP 0 011 782 A (UNION CARBIDE CORPORATION) 11 June 1980 (1980-06-11) page 10 -page 11	1,2
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

12 January 2000

Date of mailing of the international search report

28/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Rinkel, L

INTERNATIONAL SEARCH REPORT

Int. l. Application No
PCT/US 99/23794

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHUJO, YOSHIKI ET AL: "Synthesis of triethoxysilyl-terminated polyoxazolines and their cohydrolysis polymerization with tetraethoxysilane" MACROMOLECULES (1993), 26(21), 5681-6 , XP000397708 the whole document	11
X	CHEMICAL ABSTRACTS, vol. 113, no. 5, 30 July 1990 (1990-07-30) Columbus, Ohio, US; abstract no. 41014, KABETA, KEIJI: "Preparation of N,N-bis(alkoxysilylpropyl)aminohydrocarbyl amines" XP002127314 abstract & JP 02 019385 A (TOSHIBA SILICONE CO., LTD., JAPAN) 23 January 1990 (1990-01-23)	1,2

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 99/23794

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5393353	A	28-02-1995	NONE	
US 4990641	A	05-02-1991	JP 2019387 A JP 2019388 A	23-01-1990 23-01-1990
EP 11782	A	11-06-1980	US 4243767 A AU 526613 B AU 5178479 A CA 1123988 A JP 1430638 C JP 55071752 A JP 59017136 B	06-01-1981 20-01-1983 22-05-1980 18-05-1982 24-03-1988 30-05-1980 19-04-1984
JP 2019385	A	23-01-1990	NONE	